

Positron scattering resonances – dynamics and decay

J. P. Sullivan,^{1*} Z. Cheong,¹ & D. Stevens¹

*presenting author

¹james.sullivan@anu.edu.au, Australian National University

Positrons are the most easily accessible form of antimatter for use in experiments, and have been the basis of both fundamental and practical interest for many years. From a fundamental point of view, it is interesting to study the interactions of positrons with atoms and molecules, to test our understanding of these quantum interactions. This has been the focus of many research programs since the discovery of positrons in the early 1930s [1]. With the development of the Surko trap and beam system [2], a new tool was available with unprecedented energy resolution and control, opening up new avenues of investigation of low energy scattering processes. At the Australian National University, we have a beam with energy resolution of better than 50 meV [3], which is used to study a wide range of scattering processes.

One interesting problem is the question of resonance formation in positron scattering from atoms or molecules. This is a phenomenon that has been widely observed in electron scattering, with the formation of a temporary anion that will then decay and can subsequently lead to enhancement of scattering cross sections, such as vibrational excitation [4,5]. After extensive searches for such resonances in positron scattering, the first experimental evidence came with the measurement of the low energy annihilation cross section of hydrocarbons [6]. This showed a series of vibrational Feschbach resonances, which are a byproduct of the existence of a bound state between a positron and the molecule in question. Further theoretical study of these complexes predicts a coupling between the resonant state and the vibrational modes of the molecule, which should give rise to observable effects in the vibrational excitation cross sections [7]. The study presented here measures these cross sections, to elucidate the decay processes of the positron-molecule scattering resonances and compare to theoretical prediction. Positron induced vibrational excitation cross sections have been measured from a range of simple hydrocarbons - ethane, ethylene and acetylene. This talk will present this data and explore the significance of the results in terms of our understanding of positron scattering resonances.

- [1] G. F. Gribakin et al., Rev. Mod. Phys. **82**, (2010) 2577
- [2] S. J. Gilbert et al., Appl. Phys. Lett. **70**, 1944 (1997)
- [3] J. P. Sullivan et al., Rev. Sci. Instr. **79**, 113105 (2008)
- [4] S. J. Buckman and C. W. Clark, Rev. Mod. Phys. **66** (1994) 539
- [5] M J Brennan et al., J. Phys. B: At. Mol. Opt. Phys. **25** (1992) 2669
- [6] S. J. Gilbert et al., Phys. Rev. Lett. **88**, (2002) 043201
- [7] G. F. Gribakin, J. A. Young, and C. M. Surko, Rev. Mod. Phys. **82** (2010) 2557