

Stereodynamics in collisions of highly-charged ions with Ar dimers : asymmetric ion-pair formation

Tomoko Ohyama-Yamaguchi^{*1} and Atsushi Ichimura^{†2}

^{*} Tokyo Metropolitan College of Industrial Technology, Shinagawa, Tokyo 140-0011, Japan

[†] Institute of Space and Astronautical Science, JAXA, Sagami-hara, Kanagawa 252-5210, Japan

Synopsis Stereodynamical effects are analyzed for multiple ionization of rare gas dimers by slow highly charged ions using the three-center Coulombic over-barrier model previously developed by the present authors.

More than ten years ago, we proposed a three-center Coulombic over-barrier model to describe sequential multiple ionization of a rare gas dimer BC collided by a slow highly charged ion A^{q+} [1]. In more recent works [2, 3], we modified the model so as to introduce the effect of partial screening during a collision for non-active target atomic site (either B or C) and also for projectile site (A) in respective steps of electron removal. The model predicts the population distribution over charge-states (Q, Q') of dissociating ion pairs just after the collision. Measured result [4, 5] of the ion pair distribution in $Ar^{9+} + Ar_2$ collisions was reasonably reproduced in the model by taking a screening parameter as $s = 0.3 \sim 0.4$ [3].

In the present work, stereodynamical effects are examined in relation to the screening effect. We have calculated the ion pair formation cross sections as a function of the orientation angle, $\cos \theta = \hat{\mathbf{d}} \cdot \hat{\mathbf{v}}$, where \mathbf{d} denotes a molecular axis vector from C to B with \mathbf{v} being the projectile beam velocity. In addition, to obtain a physical insight more clearly, we introduce a pair of atomic impact parameters b_B and b_C respectively defined as the vertical distances from sites B and C to the incident trajectory (see Fig.1). Hence, we distinguish the *near* and *far* sites; thereby discuss the ion-pair formation cross section $\sigma(Q_{\text{near}}, Q_{\text{far}})$. The near and far sites would be determined through the measurement of momentum transfer.

Figure 2 shows the angular dependence of the cross sections $\sigma(Q_{\text{near}}, Q_{\text{far}})$ for $(Q_{\text{near}}, Q_{\text{far}}) = (2,1)$, (1,2), (3,2), and (2,3). It is seen from the figure that the (2,1) population overwhelms (1,2), and similarly (3,2) overwhelms (2,3). All the curves in the figure are symmetric with respect to $\cos \theta$. We see two maxima in (2,1) and (3,2) cross sections, and two cusps in (1,2) and (2,3) at $\cos \theta = \pm 1$. These behaviors come from the geometry of saddle point formation in the three-center Coulombic potential [2, 3].




Figure 1. Atomic impact parameters b_B for $A^{q+} + B$ and b_C for $A^{q+} + C$ with molecular impact parameter b for $A^{q+} + BC$.




Figure 2. Angular dependence of the ion-pair formation cross sections $\sigma(Q_{\text{near}}, Q_{\text{far}})$ in $A^{9+} + Ar_2$ collisions; those for (2,1) are indicated by closed circles (●), those for (1,2) by open circles (○), those for (3,2) by closed triangles (▲), and those for (2,3) by open triangles (△).

References

- [1] T. Ohyama-Yamaguchi and A. Ichimura 2003 *Nucl. Instrum. Methods B* **205** 620
- [2] T. Ohyama-Yamaguchi and A. Ichimura 2013 *Physica Scripta T* **156** 014043
- [3] T. Ohyama-Yamaguchi and A. Ichimura 2015 *J. Phys. Conf. Ser.* **635** 032101
- [4] Matsumoto *et al.* 2010 *J. Phys. Rev. Lett.* **105** 263202
- [5] Iskandar *et al.* 2014 *Phys. Rev. Lett.* **113** 143201

¹E-mail: yamaguti@metro-cit.ac.jp

²E-mail: ichimura@isas.jaxa.jp